

Stable and Strategy-Proof Matching with Flexible Allotments

Online Appendix

By JOHN WILLIAM HATFIELD, SCOTT DUKE KOMINERS, AND ALEXANDER WESTKAMP

A. THE CUMULATIVE OFFER MECHANISM

In this appendix, we formally define the cumulative offer mechanisms described in Section I.B. For any preference profile \succ , the outcome of the cumulative offer mechanism according to the strict ordering \vdash of the elements of X , denoted by $\mathcal{C}^+(\succ)$, is determined by the *cumulative offer process with respect to \vdash and \succ* as follows:

Step 0: Initialize the set of contracts *available to the hospitals (at the end of step 0)* as $A^0 = \emptyset$, and initialize the set of contracts *held by hospitals (at the end of step 0)* as $L^0 = \emptyset$.

Step $k \geq 1$: Consider the set

$$U^k \equiv \{x \in X \setminus A^{k-1} : d(x) \notin d(L^{k-1}) \text{ and } \nexists z \in (X_{d(x)} \setminus A^{k-1}) \cup \{\emptyset\} \text{ such that } z \succ_{d(x)} x\},$$

which consists of those contracts *not yet available to hospitals* that are most-preferred by doctors who do not have contracts currently held by any hospital.

- If U^k is not empty, we let y^k be the highest-ranked element of U^k according to \vdash . Doctor $d(y^k)$ *proposes* y^k , making it available to $h(y^k)$. We update the set of *available* contracts to $A^k = A^{k-1} \cup \{y^k\}$; then, the hospitals *hold* $L^k = \cup_{h \in H} C^h(A^k)$, and we proceed to step $k + 1$.
- If U^k is empty, then the cumulative offer process terminates and the outcome is given by L^{k-1} .¹

We let K denote the last proposal step of the cumulative offer process with respect to \vdash and \succ , and call A^K the set of contracts *observed* in the cumulative offer process with respect to \vdash and \succ .

Note that the sequence (y^1, \dots, y^K) of contracts proposed in the cumulative offer process with respect to \vdash and \succ is, in fact, an observable offer process (for all hospitals h). Indeed, for each proposed contract $y^k \in U^k$, we have

$$d(y^k) \notin d(L^{k-1}) = d(\cup_{h \in H} C^h(A^{k-1})) = d(\cup_{h \in H} C^h(\{y^1, \dots, y^{k-1}\})),$$

as is required for observability. Even so, however, without further assumptions on hospitals' choice functions, the outcome of a cumulative offer process need not be feasible, i.e., it might be the case that $L^K = \cup_{h \in H} C^h(A^K)$ contains more than one contract with a given doctor.

B. PROOF OF THEOREM 2

In this appendix, we prove our main result, Theorem 2.

B1. Preliminaries

We adapt the notation of Hatfield et al. (2016). For an offer process $\mathbf{x} = (x^1, \dots, x^M)$, we denote the offer process (x^1, \dots, x^m) by \mathbf{x}^m .² For an offer process $\mathbf{x} = (x^1, \dots, x^M)$, we (somewhat informally) think of \mathbf{x} as representing a sequence of proposals by doctors in a cumulative offer process (see Appendix A). With that intuition, we may think of hospital h as evaluating the set of contracts $c(\mathbf{x}^m)$ “available” at “step” m of the offer process; under a multi-division choice function with flexible allotments C^h , this implicitly involves each division $s \in S$ evaluating $c_{\rightsquigarrow s}(\mathbf{x}^m)$, the subset of $c(\mathbf{x}^m)$ that s has the opportunity to consider in the computation of

¹Note that if U^k is empty, all doctors who currently do not have a contract on hold have already proposed all the contracts they find acceptable.

²In particular, we use the convention that \mathbf{x}^0 represents the empty sequence.

$C^h(\mathbf{c}(\mathbf{x}^m))$. In the sequel, it is useful for us to track the full set of contract offers that are ever considered by s in this way by step m ; we call this set, $\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}) \equiv \cup_{m \leq M} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m)$, the set of contracts that division s has access to at some step of the offer process.

Additionally, for each division $s \in S$ we define an *extended rejection function*

$$R^s(Y; a) \equiv Y \setminus C^s(Y; a),$$

which specifies the contracts s does not choose when given a set of contract offers Y and an allotment of positions a . We note that when $C^s(\cdot; a)$ is substitutable, $R^s(\cdot; a)$ is isotone.

B2. Observable Substitutability

We show first that the choice of division s at the end of an observable offer process \mathbf{x} , i.e., $C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x})))$, coincides with the choice by s from the union of its opportunity sets along the observable offer process, i.e., $C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x})))$. Moreover, for any observable offer process \mathbf{y} such that every contract offered during \mathbf{x} is offered during \mathbf{y} (i.e., $\mathbf{c}(\mathbf{x}) \subseteq \mathbf{c}(\mathbf{y})$) we have that any contract rejected by division s when $\mathbf{c}(\mathbf{x})$ is available to h is also rejected by s when $\mathbf{c}(\mathbf{y})$ is available to h .

CLAIM 1: *If \mathbf{x} is an observable offer process, then for each division $s \in S$,*

$$(B1) \quad C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x}))) = C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x}))).$$

Furthermore, if \mathbf{y} is an observable offer process such that $\mathbf{c}(\mathbf{x}) \subseteq \mathbf{c}(\mathbf{y})$, and $\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}) \subseteq \mathbf{f}_{\rightsquigarrow s}(\mathbf{y})$ for some division $s \in S$, then

$$(B2) \quad R^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x}))) \subseteq R^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{y}); q^s(\mathbf{c}(\mathbf{y}))).$$

PROOF:

We fix an observable offer process $\mathbf{x} = (x^1, \dots, x^M)$. We proceed by induction on pairs (m, s) in the order

$$(1, 1), (1, 2), \dots, (1, \bar{s}), (2, 1), (2, 2), \dots, (2, \bar{s}), \dots, (M, 1), (M, 2), \dots, (M, \bar{s}),$$

showing at each step that

$$(B3a) \quad C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))) = C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))),$$

$$(B3b) \quad C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))) \subseteq C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) \cup [\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1})],$$

$$(B3c) \quad C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) \subseteq \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m),$$

$$(B3d) \quad R^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))) \subseteq R^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{y}); q^s(\mathbf{c}(\mathbf{y})))$$

for all observable offer processes \mathbf{y} such that $\mathbf{c}(\mathbf{x}^m) \subseteq \mathbf{c}(\mathbf{y})$ and divisions s such that $\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m) \subseteq \mathbf{f}_{\rightsquigarrow s}(\mathbf{y})$. Taking $m = M$ then provides the desired results via (B3a) and (B3d).

Base Case(s). For $m = 1$, conditions (B3a), (B3b), and (B3c) follow immediately for all $s \in S$.

Condition (B3d) follows using the same argument as we use in the general (m, s) case *infra*.

Inductive Step. We now show that (B3) holds for (m, s) if (B3) holds for

- every pair (\hat{m}, t) with $\hat{m} < m$ and $t \in S$, as well as
- every pair (m, t) with $t < s$.

Condition (B3c). First, we note that the result is immediate if $s = 1$, as $\mathbf{c}_{\rightsquigarrow 1}(\mathbf{x}^{m-1}) = \mathbf{c}(\mathbf{x}^{m-1}) \subseteq \mathbf{c}(\mathbf{x}^m) = \mathbf{c}_{\rightsquigarrow 1}(\mathbf{x}^m)$. Thus, we consider any contract z chosen by division $s > 1$ at step $m - 1$,

that is, any $z \in C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1})))$. Under the choice procedure defining C^h , since $z \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1})$, it must be that no contract with $d(z)$ is chosen by any division $t < s$ at step $m - 1$, i.e.,

$$[\mathbf{c}(\mathbf{x}^{m-1})]_{d(z)} \subseteq \bigcap_{t < s} R^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(\mathbf{c}(\mathbf{x}^{m-1}))).$$

Moreover, conditions (B3a) and (B3b) together imply that for each $t < s$,

$$C^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^m); q^t(\mathbf{c}(\mathbf{x}^m))) \subseteq C^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(\mathbf{c}(\mathbf{x}^{m-1}))) \cup [\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^m) \setminus \mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{m-1})].$$

Hence, we have

$$(B4) \quad [\mathbf{c}(\mathbf{x}^{m-1})]_{d(z)} \subseteq \bigcap_{t < s} R^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^m); q^t(\mathbf{c}(\mathbf{x}^m))),$$

i.e., it must be that no contract with $d(z)$ —except possibly x^m —is chosen by any division $t < s$ at step m . But since \mathbf{x}^m is observable and $z \in C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1})))$, it must be that $d(z) \neq d(x^m)$. Thus, (B4) implies

$$[\mathbf{c}(\mathbf{x}^m)]_{d(z)} \subseteq \bigcap_{t < s} R^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^m); q^t(\mathbf{c}(\mathbf{x}^m))),$$

that is, no contract with $d(z)$ is chosen by any division $t < s$ at step m . Thus, any contract proposed by $d(z)$ —and, in particular, z —is available to s at step m . Therefore, since z was an arbitrary element of $C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1})))$, we have that $C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) \subseteq \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m)$, as desired.

Condition (B3b). Taking $\mathbf{y} = \mathbf{x}^m$ in condition (B3d) for $(m - 1, s)$, we obtain

$$R^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) \subseteq R^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))),$$

which implies that

$$C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))) \subseteq C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) \cup [\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus \mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^{m-1})].$$

Since $\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m) = \bigcup_{n \leq m} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)$ and $\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^{m-1}) = \bigcup_{n \leq m-1} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)$ by definition, we have

$$\begin{aligned} \mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus \mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^{m-1}) &= [\bigcup_{n \leq m} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)] \setminus [\bigcup_{n \leq m-1} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)] \\ &= [\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus [\bigcup_{n \leq m-1} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)]] \cup [[\bigcup_{n \leq m-1} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)] \setminus [\bigcup_{n \leq m-1} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)]] \\ &= [\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus [\bigcup_{n \leq m-1} \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^n)]] \cup \emptyset \\ &\subseteq \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}). \end{aligned}$$

Combining the two immediately preceding expressions yields

$$(B5) \quad C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))) \subseteq C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) \cup [\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1})].$$

Now, since condition (B3a) holds for $(m - 1, s)$, we have that

$$(B6) \quad C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) = C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))).$$

Combining (B5) and (B6) implies that

$$C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))) \subseteq C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(\mathbf{c}(\mathbf{x}^{m-1}))) \cup [\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^m) \setminus \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{m-1})],$$

as desired.

Condition (B3a). Condition (B3b) for (m, s) implies that

$$C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \subseteq C^s(c_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(c(\mathbf{x}^{m-1}))) \cup [c_{\rightsquigarrow s}(\mathbf{x}^m) \setminus c_{\rightsquigarrow s}(\mathbf{x}^{m-1})];$$

condition (B3c) implies that $C^s(c_{\rightsquigarrow s}(\mathbf{x}^{m-1}); q^s(c(\mathbf{x}^{m-1}))) \subseteq c_{\rightsquigarrow s}(\mathbf{x}^m)$, and so

$$C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \subseteq c_{\rightsquigarrow s}(\mathbf{x}^m).$$

Since $f_{\rightsquigarrow s}(\mathbf{x}^m) \supseteq c_{\rightsquigarrow s}(\mathbf{x}^m)$, the fact that the extended choice function of division s satisfies the irrelevance of rejected contracts condition then implies that

$$C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) = C^s(c_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))),$$

as desired.

Condition (B3d). Suppose that \mathbf{y} is an observable offer process such that $c(\mathbf{x}^m) \subseteq c(\mathbf{y})$ and $f_{\rightsquigarrow s}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow s}(\mathbf{y})$ for some $s \in S$. There are two cases to consider:

Case 1: $q^s(c(\mathbf{x}^m)) \geq q^s(c(\mathbf{y}))$. Since $f_{\rightsquigarrow s}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow s}(\mathbf{y})$, the substitutability of the extended choice function of division s implies that $R^s(c(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \subseteq R^s(c(\mathbf{y}); q^s(c(\mathbf{x}^m)))$. Since $q^s(c(\mathbf{x}^m)) \geq q^s(c(\mathbf{y}))$ and the extended choice function of s is monotonic with respect to the allotment, we have $R^s(c(\mathbf{y}); q^s(c(\mathbf{x}^m))) \subseteq R^s(c(\mathbf{y}); q^s(c(\mathbf{y})))$ and so

$$(B7) \quad R^s(c(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \subseteq R^s(c(\mathbf{y}); q^s(c(\mathbf{y}))),$$

as desired.

Case 2: $q^s(c(\mathbf{x}^m)) < q^s(c(\mathbf{y}))$. We first show that

$$(B8) \quad C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) = C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty).$$

To show (B8), we suppose the contrary—i.e., that $C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \neq C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$ —and seek a contradiction. Now, if $C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \neq C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$, then, as the extended choice function of s is monotonic with respect to the allotment, we must have $C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \subsetneq C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$. We let $z \in C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty) \setminus C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m)))$. As $z \in f_{\rightsquigarrow s}(\mathbf{x}^m)$, there must exist some largest $\bar{m} \leq m$ such that $z \in c_{\rightsquigarrow s}(\mathbf{x}^{\bar{m}})$. There are then two subcases to consider:

If $\bar{m} = m$, then since condition (B3a) holds for (m, s) , we must have that $C^s(c_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) = C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m)))$. As $z \notin C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m)))$ and $z \in c_{\rightsquigarrow s}(\mathbf{x}^m)$, we then have that

$$z \in R^s(c_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))).$$

As the allotment function is single-peaked across observable offer processes and $q^s(c(\mathbf{x}^m)) < q^s(c(\mathbf{y}))$, we have that $R^s(c_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) = R^s(c_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$; in particular, $z \in R^s(c_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$. Thus, as the extended choice function of s is substitutable, we have $z \in R^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$, contradicting our assumption that $z \in C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$.

If $\bar{m} < m$, then $z \notin c_{\rightsquigarrow s}(\mathbf{x}^{\bar{m}+1})$, as we chose \bar{m} to be the largest $\bar{m} \leq m$ such that $z \in c_{\rightsquigarrow s}(\mathbf{x}^{\bar{m}})$. Thus, $z \in R^s(c_{\rightsquigarrow s}(\mathbf{x}^{\bar{m}}); q^s(c(\mathbf{x}^{\bar{m}})))$, as otherwise condition (B3c) is violated. As the allotment function is single-peaked across observable offer processes and $q^s(c(\mathbf{x}^m)) < q^s(c(\mathbf{y}))$, we have that $R^s(c_{\rightsquigarrow s}(\mathbf{x}^{\bar{m}}); q^s(c(\mathbf{x}^{\bar{m}}))) = R^s(c_{\rightsquigarrow s}(\mathbf{x}^{\bar{m}}); \infty)$; in particular, $z \in R^s(c_{\rightsquigarrow s}(\mathbf{x}^{\bar{m}}); \infty)$. Thus, as the extended choice function of s is

substitutable, $z \in R^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$. Then, again as the extended choice function of s is substitutable, we must have $z \in R^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$, contradicting our assumption that $z \in C^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$.

The preceding argument shows (B8), which implies that $R^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(\mathbf{c}(\mathbf{x}^m))) = R^s(f_{\rightsquigarrow s}(\mathbf{x}^m); \infty)$. The substitutability of the extended choice function of s then implies that $R^s(f_{\rightsquigarrow s}(\mathbf{c}(\mathbf{x}^m)); q^s(\infty)) \subseteq R^s(f_{\rightsquigarrow s}(\mathbf{y}); \infty)$, and so

$$R^s(f_{\rightsquigarrow s}(\mathbf{c}(\mathbf{x}^m)); q^s(\mathbf{c}(\mathbf{x}^m))) \subseteq R^s(f_{\rightsquigarrow s}(\mathbf{y}); \infty).$$

Finally, the monotonicity of the extended choice function of s with respect to allotment (i.e., $R^s(f_{\rightsquigarrow s}(\mathbf{y}); \infty) \subseteq R^s(f_{\rightsquigarrow s}(\mathbf{y}); q^s(\mathbf{c}(\mathbf{y})))$) implies (B7) in this case too, completing our induction.

We now show that the choice function of h is observably substitutable.

CLAIM 2: *The choice function C^h is observably substitutable.*

PROOF:

We consider an observable offer process $\mathbf{x} = (x^1, \dots, x^M)$ and let $y \in R^h(\{x^1, \dots, x^{M-1}\})$. Under the choice procedure defining C^h , since $y \in R^h(\{x^1, \dots, x^{M-1}\})$, we have that $y \in R^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1})))$ for each $s \in S$. The substitutability of the extended choice function of each $s \in S$ then implies that $y \in R^s(f_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1})))$ for each $s \in S$. Since $f_{\rightsquigarrow s}(\mathbf{x}^{M-1}) \subseteq f_{\rightsquigarrow s}(\mathbf{x}^M)$ by construction, (B2) of Claim 1 implies that $y \in R^s(f_{\rightsquigarrow s}(\mathbf{x}^M); q^s(\mathbf{c}(\mathbf{x}^M)))$ for each $s \in S$. Thus, $y \notin C^s(f_{\rightsquigarrow s}(\mathbf{x}^M); q^s(\mathbf{c}(\mathbf{x}^M)))$ for each $s \in S$; (B1) of Claim 1 then implies that $y \notin C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^M); q^s(\mathbf{c}(\mathbf{x}^M)))$ for each $s \in S$. Thus, under the choice procedure defining C^h , we have that $y \in R^h(\{x^1, \dots, x^M\})$, as desired.

B3. Observable Size Monotonicity

Next, we show that the choice function C^h is observably size monotonic across observable offer processes.

CLAIM 3: *The choice function C^h is observably size monotonic.*

PROOF:

Consider any two observable offer process \mathbf{x} and $\hat{\mathbf{x}}$ such that $\mathbf{c}(\mathbf{x}) \subseteq \mathbf{c}(\hat{\mathbf{x}})$. As the allotment function does not observably grant excess positions, we have that $q^s(\mathbf{c}(\mathbf{x})) = |C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x})))|$ for each division $s \in S$, which implies that $\sum_{s \in S} q^s(\mathbf{c}(\mathbf{x})) = \sum_{s \in S} |C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x})))|$. Similarly, we have that $\sum_{s \in S} q^s(\mathbf{c}(\hat{\mathbf{x}})) = \sum_{s \in S} |C^s(\mathbf{c}_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(\mathbf{c}(\hat{\mathbf{x}})))|$. Now, as the allotment function is monotone in aggregate across observable offer processes, we have that

$$\sum_{s \in S} q^s(\mathbf{c}(\mathbf{x})) \leq \sum_{s \in S} q^s(\mathbf{c}(\hat{\mathbf{x}}));$$

hence, we have

$$|C^h(\mathbf{c}(\mathbf{x}))| = \sum_{s \in S} |C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x})))| \leq \sum_{s \in S} |C^s(\mathbf{c}_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(\mathbf{c}(\hat{\mathbf{x}})))| = |C^h(\mathbf{c}(\hat{\mathbf{x}}))|,$$

so the choice function C^h is observably size monotonic, as desired.

B4. (Non-)Manipulability via Contractual Terms

We now establish that C^h is non-manipulable via contractual terms. Consider an arbitrary doctor $d \in D$, and let z^0, z^1, \dots, z^N be an arbitrary sequence of contracts in X_d . Fix a profile of

preferences $\succ_{D \setminus \{d\}}$ for all other doctors, and let \succ_d and $\hat{\succ}_d$ be given by

$$(B9) \quad \succ_d : z^1 \succ_d \dots \succ_d z^N,$$

$$(B10) \quad \hat{\succ}_d : z^0 \hat{\succ}_d z^1 \hat{\succ}_d \dots \hat{\succ}_d z^N.$$

We fix an ordering \vdash over the set of contracts X , and let $\mathbf{x} = (x^1, \dots, x^M)$ be the observable offer process induced by the cumulative offer mechanism with ordering \vdash under the preferences $(\succ_d, \succ_{D \setminus \{d\}})$ when only hospital h is present. Similarly, let $\hat{\mathbf{x}} = (\hat{x}^1, \dots, \hat{x}^M)$ be the observable offer process induced by the cumulative offer mechanism with ordering \vdash under the preferences $(\hat{\succ}_d, \succ_{D \setminus \{d\}})$ when only hospital h is present. We first establish the following claim.

CLAIM 4: *If $z^0 \notin C^h(c(\hat{\mathbf{x}}))$, then*

$$(B11) \quad R^h(c(\mathbf{x})) \subseteq R^h(c(\hat{\mathbf{x}}))$$

and, for all $s \in S$, we have that

$$(B12) \quad f_{\rightsquigarrow s}(\mathbf{x}) \subseteq f_{\rightsquigarrow s}(\hat{\mathbf{x}}).$$

PROOF:

We proceed by induction on pairs (m, s) in the order

$$(1, 1), (1, 2), \dots, (1, \bar{s}), (2, 1), (2, 2), \dots, (2, \bar{s}), \dots, (M, 1), (M, 2), \dots, (M, \bar{s}),$$

showing at each step that

$$(B13a) \quad f_{\rightsquigarrow s}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow s}(\hat{\mathbf{x}}),$$

$$(B13b) \quad R^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \subseteq R^s(f_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(c(\hat{\mathbf{x}}))).$$

Once we have completed our induction, (B12) follows from taking $m = M$.

For the base case $(1, 1)$, it must be the case that x^1 is either the highest-ranked contract by some doctor $d(x^1) \neq d$ or $x^1 = z^1$. In the former case, x^1 must be offered at some step in the offer process $\hat{\mathbf{x}}$, as it is the favored contract of doctor $d(x^1)$. In the latter case, since z^0 is rejected by h by assumption, d must offer his second-favorite contract under $\hat{\succ}_d$, i.e., x^1 , at some step in the offer process $\hat{\mathbf{x}}$. Hence, in both cases we have that $x^1 \in c(\hat{\mathbf{x}})$. Since $f_{\rightsquigarrow 1}(\hat{\mathbf{x}}) = c(\hat{\mathbf{x}})$, we obtain that $f_{\rightsquigarrow 1}(\mathbf{x}^1) \subseteq f_{\rightsquigarrow 1}(\hat{\mathbf{x}})$. Then, condition (B2) of Claim 1 implies that $R^1(f_{\rightsquigarrow 1}(\mathbf{x}^1); q^1(c(\mathbf{x}^1))) \subseteq R^1(f_{\rightsquigarrow 1}(\hat{\mathbf{x}}); q^1(c(\hat{\mathbf{x}})))$.

We now show that (B13) holds for (m, s) if (B13) holds for

- every pair (\bar{m}, t) with $\bar{m} < m$ and $t \in S$, as well as
- every pair (m, t) with $t < s$.

We show first that (B13) holds for $(m, 1)$ given that (B13) holds for every pair (\bar{m}, t) with $\bar{m} < m$ and $t \in S$. By the inductive assumption (B13b) for pairs $(m-1, t)$, we have

$$R^t(f_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(c(\mathbf{x}^{m-1}))) \subseteq R^t(f_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}})))$$

for all t . As \mathbf{x}^m is observable, we have that $\{x^1, \dots, x^{m-1}\}_{d(x^m)} \subseteq R^t(c_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(c(\mathbf{x}^{m-1})))$ for all t . Moreover, as $R^t(\cdot; q^t(c(\mathbf{x}^{m-1})))$ is substitutable, we have $R^t(c_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(c(\mathbf{x}^{m-1}))) \subseteq R^t(f_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(c(\mathbf{x}^{m-1})))$, so that

$$\{x^1, \dots, x^{m-1}\}_{d(x^m)} \subseteq R^t(c_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(c(\mathbf{x}^{m-1}))) \subseteq R^t(f_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(c(\mathbf{x}^{m-1}))).$$

Hence, given that $R^t(f_{\rightsquigarrow t}(\mathbf{x}^{m-1}); q^t(c(\mathbf{x}^{m-1}))) \subseteq R^t(f_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}})))$ for all t , we find that $\{x^1, \dots, x^{m-1}\}_{d(x^m)} \subseteq R^t(f_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}})))$ for all t .

By Condition (B1) of Claim 1, we see that there is an $M' \leq \hat{M}$ such that $\{x^1, \dots, x^{m-1}\}_{d(x^m)} \subseteq R^t(c_{\rightsquigarrow t}(\hat{\mathbf{x}}^{M'}); q^t(c(\hat{\mathbf{x}}^{M'})))$ for all t . Since $\hat{\mathbf{x}}$ is observable and represents all the offers made under the cumulative offer process for $(\succ_d, \succ_{D \setminus \{d\}})$, there must exist some step \hat{m} at which x^m is proposed in $\hat{\mathbf{x}}$; hence, recalling that $c(\hat{\mathbf{x}}) = f_{\rightsquigarrow 1}(\hat{\mathbf{x}})$, we have that

$$(B14) \quad x^m \in c(\hat{\mathbf{x}}) = f_{\rightsquigarrow 1}(\hat{\mathbf{x}}).$$

Additionally, by the inductive assumption (B13a) for $m-1$ and $s=1$, we have that

$$(B15) \quad c(\mathbf{x}^{m-1}) = f_{\rightsquigarrow 1}(\mathbf{x}^{m-1}) \subseteq f_{\rightsquigarrow 1}(\hat{\mathbf{x}}).$$

Moreover, recalling that $f_{\rightsquigarrow 1}(\mathbf{x}^m) = c(\mathbf{x}^m)$ and $c(\mathbf{x}^m) = \{x^m\} \cup c(\mathbf{x}^{m-1})$, we have that

$$(B16) \quad f_{\rightsquigarrow 1}(\mathbf{x}^m) = c(\mathbf{x}^m) = \{x^m\} \cup c(\mathbf{x}^{m-1}).$$

Combining (B14) and (B15) with (B16) then implies that $f_{\rightsquigarrow 1}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow 1}(\hat{\mathbf{x}})$, which is exactly condition (B13a) for $(m, 1)$. Applying condition (B2) of Claim 1 yields that $R^1(f_{\rightsquigarrow 1}(\mathbf{x}^m); q^1(c(\mathbf{x}^m))) \subseteq R^1(f_{\rightsquigarrow 1}(\hat{\mathbf{x}}); q^1(c(\hat{\mathbf{x}})))$, i.e., condition (B13b) for $(m, 1)$.

We now show that (B13) holds for (m, s) when $s > 1$, given that (B13) holds for every pair (\bar{m}, t) with $\bar{m} < m$ and $t \in S$ and every pair (m, t) with $t < s$. We argue first that $f_{\rightsquigarrow s}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow s}(\hat{\mathbf{x}})$. By our inductive assumption on $(m-1, s)$, it is sufficient to show that $c_{\rightsquigarrow s}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow s}(\hat{\mathbf{x}})$, as $f_{\rightsquigarrow s}(\mathbf{x}^m) = f_{\rightsquigarrow s}(\mathbf{x}^{m-1}) \cup c_{\rightsquigarrow s}(\mathbf{x}^m)$. Let $y \in c_{\rightsquigarrow s}(\mathbf{x}^m)$ be arbitrary. Since $y \in c_{\rightsquigarrow s}(\mathbf{x}^m)$, under the choice procedure defining C^h , we must have that

$$(B17) \quad y \in \bigcap_{t < s} R^t(c_{\rightsquigarrow t}(\mathbf{x}^m); q^t(c(\mathbf{x}^m))).$$

As each extended choice function C^t is substitutable, (B17) implies that

$$(B18) \quad y \in \bigcap_{t < s} R^t(f_{\rightsquigarrow t}(\mathbf{x}^m); q^t(c(\mathbf{x}^m))).$$

Now, by the inductive assumption (B13b) on pairs (m, t) for $t < s$, we have that $R^t(f_{\rightsquigarrow t}(\mathbf{x}^m); q^t(c(\mathbf{x}^m))) \subseteq R^t(f_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}})))$ for all $t < s$. Hence, (B18) implies that

$$(B19) \quad y \in \bigcap_{t < s} R^t(f_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}}))).$$

Applying condition (B1) of Claim 1 to $\hat{\mathbf{x}}$, we have that $C^t(c_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}}))) = C^t(f_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}})))$ for all t . Thus, if there were a $t < s$ such that $y \in C^t(c_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}})))$, we would have $y \in C^t(f_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}})))$, contradicting (B19). Hence, we must have

$$y \in \bigcap_{t < s} R^t(c_{\rightsquigarrow t}(\hat{\mathbf{x}}); q^t(c(\hat{\mathbf{x}}))).$$

Thus, there must exist some step \hat{m} of the offer process $\hat{\mathbf{x}}$ such that $y \in c_{\rightsquigarrow s}(\hat{\mathbf{x}}^{\hat{m}})$, and so $y \in f_{\rightsquigarrow s}(\hat{\mathbf{x}}^{\hat{m}}) \subseteq f_{\rightsquigarrow s}(\hat{\mathbf{x}})$; hence, we see that $f_{\rightsquigarrow s}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow s}(\hat{\mathbf{x}})$, i.e., we have condition (B13a) for (m, s) . Applying condition (B2) of Claim 1 then gives that $R^s(f_{\rightsquigarrow s}(\mathbf{x}^m); q^s(c(\mathbf{x}^m))) \subseteq R^s(f_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(c(\hat{\mathbf{x}})))$ —condition (B13b) for (m, s) . Having thus completed our induction, we have proven (B12).

Finally, we prove (B11), i.e., that $R^h(\mathbf{c}(\mathbf{x})) \subseteq R^h(\mathbf{c}(\hat{\mathbf{x}}))$. Suppose by way of contradiction that there exists some $y \in R^h(\mathbf{c}(\mathbf{x})) \setminus R^h(\mathbf{c}(\hat{\mathbf{x}}))$. We have already shown (B12), that $f_{\rightsquigarrow s}(\mathbf{x}) \subseteq f_{\rightsquigarrow s}(\hat{\mathbf{x}})$ for all $s \in S$; in particular,

$$y \in \mathbf{c}(\mathbf{x}) = f_{\rightsquigarrow 1}(\mathbf{x}^m) \subseteq f_{\rightsquigarrow 1}(\hat{\mathbf{x}}) = \mathbf{c}(\hat{\mathbf{x}}).$$

Thus, as $y \notin R^h(\mathbf{c}(\hat{\mathbf{x}}))$, we must have that $y \in C^h(\mathbf{c}(\hat{\mathbf{x}}))$. Therefore, there exists some division $s \in S$ such that $y \in C^s(\mathbf{c}_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(\hat{\mathbf{x}}))$; by (B1) of Claim 1, we have that $y \in C^s(f_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(\hat{\mathbf{x}}))$ and so $y \notin R^s(f_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(\hat{\mathbf{x}}))$. By (B2) of Claim 1, since $\mathbf{c}(\mathbf{x}) \subseteq \mathbf{c}(\hat{\mathbf{x}})$, we then have that $y \notin R^s(f_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{x}))$. But as the extended choice function of s is substitutable, we must then have that $y \notin R^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{x}))$ —but this can only be the case if s or some division $t < s$ chooses y , which would contradict the assumption that $y \in R^h(\mathbf{c}(\mathbf{x}))$.

CLAIM 5: *The choice function C^h is not manipulable via contractual terms.*

PROOF:

By Proposition 5 of Hatfield et al. (2016), it is sufficient to show that when h is the only hospital, the following two conditions hold:

- 1) If $[\mathcal{C}(\succ_d, \succ_{D \setminus \{d\}})]_d = \emptyset$, then either $[\mathcal{C}(\hat{\succ}_d, \succ_{D \setminus \{d\}})]_d = \emptyset$ or $[\mathcal{C}(\hat{\succ}_d, \succ_{D \setminus \{d\}})]_d = \{z^0\}$, and
- 2) if $[\mathcal{C}(\hat{\succ}_d, \succ_{D \setminus \{d\}})]_d = \emptyset$, then $[\mathcal{C}(\succ_d, \succ_{D \setminus \{d\}})]_d = \emptyset$.³

To show the first condition, we note that Claim 4 implies that if $[\mathcal{C}(\succ_d, \succ_{D \setminus \{d\}})]_d = \emptyset$ and $z^0 \notin \mathcal{C}(\hat{\succ}_d, \succ_{D \setminus \{d\}})$, then $R^h(\mathbf{c}(\mathbf{x})) \subseteq R^h(\mathbf{c}(\hat{\mathbf{x}}))$. Moreover, if $[\mathcal{C}(\succ_d, \succ_{D \setminus \{d\}})]_d = \emptyset$, then $\{z^1, \dots, z^N\} \subseteq R^h(\mathbf{c}(\mathbf{x}))$, and so combining the preceding two observations, we have that $\{z^1, \dots, z^N\} \subseteq R^h(\mathbf{c}(\hat{\mathbf{x}}))$; hence $[\mathcal{C}(\hat{\succ}_d, \succ_{D \setminus \{d\}})]_d = \emptyset$.

To show the second condition, note that by Proposition 1 of Hatfield et al. (2016), since the choice function of h is observably substitutable (Claim 2) and observably size monotonic (Claim 3), the cumulative offer mechanism outcome is order-independent. Thus we can consider \mathbf{x} and $\hat{\mathbf{x}}$ to be generated by cumulative offer processes with respect to the proposal ordering \vdash in which all of the contracts associated with doctors other than d precede all of the contracts associated with d , i.e., if $x \in X_d$ and $y \in X_{D \setminus \{d\}}$, then $y \vdash x$.

Under our choice of \vdash , there must exist an \bar{m} such that

- 1) $x^m = \hat{x}^m$ for all $m < \bar{m}$,
- 2) $x^{\bar{m}} = z^1$, and
- 3) $\hat{x}^{\bar{m}} = z^0$;

specifically, \bar{m} is the first step of each cumulative offer process (with respect to \vdash) at which d proposes. Additionally, at each step after \bar{m} , exactly one contract is newly rejected and the offer process $\hat{\mathbf{x}}$ must end with the contract z^N being rejected, as

- z^N follows all contracts with doctors other than d under \vdash ,
- the choice function of h is observably substitutable,
- the choice function of h is observably size monotonic, and
- we have assumed that $[\mathcal{C}(\hat{\succ}_d, \succ_{D \setminus \{d\}})]_d = \emptyset$.

Thus, we must have that

- 1) $|R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}})) \setminus R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}-1}))| = 1$ for all $\hat{m} \in \{\bar{m}, \bar{m} + 1, \dots, \hat{M}\}$, and

³Here, we use the notation introduced in (B9).

2) $z^N \in R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{M}})) \setminus R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{M}-1}))$.

Similarly, for the offer process \mathbf{x} , we must have $|R^h(\mathbf{c}(\mathbf{x}^m)) \setminus R^h(\mathbf{c}(\mathbf{x}^{m-1}))| = 1$ for all $m \in \{\bar{m}, \bar{m} + 1, \dots, M - 1\}$.

Since $\mathbf{x}^{\bar{m}-1} = \hat{\mathbf{x}}^{\bar{m}-1}$, we have that $|C^h(\mathbf{c}(\mathbf{x}^{\bar{m}-1}))| = |C^h(\mathbf{c}(\hat{\mathbf{x}}^{\bar{m}-1}))|$. Moreover, since $|R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}})) \setminus R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}-1}))| = 1$ for all $\hat{m} \in \{\bar{m}, \bar{m} + 1, \dots, \hat{M}\}$ (as we have just shown), we have that $|C^h(\mathbf{c}(\mathbf{x}^{\bar{m}-1}))| = |C^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{M}}))|$. Likewise, since $|R^h(\mathbf{c}(\mathbf{x}^m)) \setminus R^h(\mathbf{c}(\mathbf{x}^{m-1}))| = 1$ for $m \in \{\bar{m}, \bar{m} + 1, \dots, M - 1\}$ (as we have just shown), we have that $|C^h(\mathbf{c}(\mathbf{x}^{M-1}))| = |C^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{M}}))|$. But since $\mathbf{c}(\mathbf{x}^M) \subseteq \mathbf{c}(\hat{\mathbf{x}}^{\hat{M}})$, the observable size monotonicity of C^h implies that $|C^h(\mathbf{c}(\mathbf{x}^M))| \leq |C^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{M}}))|$. Thus, we must have

$$(B20) \quad R^h(\mathbf{c}(\mathbf{x}^M)) \setminus R^h(\mathbf{c}(\mathbf{x}^{M-1})) \neq \emptyset.$$

Now, given (B20), suppose by way of contradiction that $y \in R^h(\mathbf{c}(\mathbf{x}^M)) \setminus R^h(\mathbf{c}(\mathbf{x}^{M-1})) \neq \emptyset$ is not the contract z^N , and so is the least preferred acceptable contract with respect to $\succ_{d(y)}$, where $d(y) \neq d$. Claim 4 then implies there is some step $\hat{m} \geq \bar{m}$ such that $y \in R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}})) \setminus R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}-1}))$. But, since $|R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}})) \setminus R^h(\mathbf{c}(\hat{\mathbf{x}}^{\hat{m}-1}))| = 1$, and y is the least preferred acceptable contract for $d(y)$, the cumulative offer process for $(\succ_d, \succ_{D \setminus \{d\}})$ would end at \hat{m} with the rejection of y , contradicting the fact that it ends with the rejection of z^N .

B5. Irrelevance of Rejected Contracts

Finally, we show that the choice function C^h satisfies the irrelevance of rejected contracts condition.

CLAIM 6: *The choice function C^h satisfies the irrelevance of rejected contracts condition.*

PROOF:

We suppose that z and Y are such that $z \in R^h(Y \cup \{z\})$. If $z \in R^h(Y \cup \{z\})$, then we must have

$$(B21) \quad z \notin C^s([Y \cup \{z\}]_{\rightsquigarrow s}; q^s(Y \cup \{z\}))$$

for each division s . Moreover, since each extended choice function C^s satisfies the irrelevance of rejected contracts condition, (B21) implies that

$$(B22) \quad C^s([Y \cup \{z\}]_{\rightsquigarrow s}; q^s(Y \cup \{z\})) = C^s(Y_{\rightsquigarrow s}; q^s(Y \cup \{z\}))$$

for all divisions s .

As the allotment function does not depend on irrelevant contracts, we have that $q(Y \cup \{z\}) = q(Y)$. Thus, we have

$$(B23) \quad C^s([Y \cup \{z\}]_{\rightsquigarrow s}; q^s(Y \cup \{z\})) = C^s([Y \cup \{z\}]_{\rightsquigarrow s}; q^s(Y)).$$

Combining (B22) and (B23) shows that

$$C^s([Y \cup \{z\}]_{\rightsquigarrow s}; q^s(Y \cup \{z\})) = C^s(Y_{\rightsquigarrow s}; q^s(Y))$$

for all divisions s ; it follows that $C^h(Y \cup \{z\}) = C^h(Y)$.

C. A MULTI-DIVISION CHOICE FUNCTION WITH FLEXIBLE ALLOTMENTS THAT IS NOT SUBSTITUTABLY COMPLETABLE

In many application contexts with non-substitutable preferences, stable and strategy-proof matching can be guaranteed by showing that each hospital's choice function is *substitutably*

completable in the sense of Hatfield and Kominers (2016). However, our main result (Theorem 2) can not be demonstrated by using substitutable completability arguments; to prove this, we show in this appendix that there is a multi-division choice function with flexible allotments that is not substitutably completable.

Specifically, we give a multi-division choice function with flexible allotments that expresses the preferences introduced in Example 2 of Hatfield et al. (2016). First, we recall the setting: $H = \{h\}$, $D = \{d, e, f\}$, and $X = \{x, y, z, \hat{x}, \hat{z}\}$, with $h(x) = h(y) = h(z) = h(\hat{x}) = h(\hat{z}) = h$, $d(x) = d(\hat{x}) = d$, $d(y) = e$, and $d(z) = d(\hat{z}) = f$. We let the choice function C^h of h be induced by the preference relation⁴

$$(C1) \quad \{\hat{x}, z\} \succ \{x, \hat{z}\} \succ \{y, \hat{z}\} \succ \{\hat{x}, y\} \succ \{x, y\} \succ \{y, z\} \succ \{\hat{x}, \hat{z}\} \succ \{x, z\} \\ \succ \{y\} \succ \{\hat{z}\} \succ \{\hat{x}\} \succ \{x\} \succ \{z\} \succ \emptyset.$$

Hatfield et al. (2016) proved that C^h does not have a substitutable completion. Here, we show that C^h can be modeled as a multi-division choice function with flexible allotments. We let $S = \{1, 2, 3, 4\}$, with the extended choice functions of the divisions s induced by the following preference relations:⁵

$$\succ_1 : \{\hat{x}\} \succ \emptyset, \\ \succ_2 : \{x, y\} \succ \{x\} \succ \{y\} \succ \emptyset, \\ \succ_3 : \{\hat{z}\} \succ \emptyset, \\ \succ_4 : \{z\} \succ \emptyset.$$

It is immediate that for each $s \in S$, the extended choice function C^s is substitutable and size monotonic, satisfies the irrelevance of rejected contracts condition for any allotment, and, moreover, is monotonic with respect to the allotment and conditionally acceptant.

For each set of contracts $Y \subseteq X$, we denote the allotment function by

$$q(Y) = (q_1(Y), q_2(Y), q_3(Y), q_4(Y));$$

in Table C1, we define the allotment function for every possible set of contracts available to h , and also state the choice of h from that set of contracts.

It is clear from Table C1 that the choice function just defined is equivalent to the choice function induced by (C1). Moreover, it is straightforward to check using Table C1 that q does not depend on irrelevant contracts, does not observably grant excess positions, and is monotone in aggregate across observable offer processes.

We now show that q is single-peaked across observable offer processes. We say that a division s is *capacity constrained under Y at n* if $C^s(Y_{\rightsquigarrow s}; q^s(Y)) \subsetneq C^s(Y_{\rightsquigarrow s}; \infty)$ and $|C^s(Y_{\rightsquigarrow s}; q^s(Y))| = n$. Thus, to show that q is single-peaked across observable offer processes, it suffices to show that if a division is capacity constrained under $c(\mathbf{x})$ at n for some offer process \mathbf{x} , for any offer process \mathbf{y} such that $c(\mathbf{x}) \subseteq c(\mathbf{y})$, division s is capacity constrained under $c(\mathbf{y})$ at $m \leq n$. Now, we consider any observable offer processes \mathbf{x} and \mathbf{y} such that $c(\mathbf{x}) \subsetneq c(\mathbf{y})$. If $|c(\mathbf{x})| \leq 2$, then no division is

⁴A preference relation \succ_h for hospital h induces a choice function C^h for h under which

$$C^h(Y) = \max_{\succ_h} \{Z \subseteq X_h : Z \subseteq Y\},$$

where by \max_{\succ_h} we mean the maximum with respect to the ordering \succ_h ; that is, h chooses its most-preferred subset of Y .

⁵A preference relation \succ_s for s induces an extended choice function C^s for s , under which

$$C^s(Y; a) = \max_{\succ_s} \{Z \subseteq Y : |Z| \leq a\}$$

where by \max_{\succ_s} we mean the maximum with respect to the ordering \succ_s ; that is, s chooses its most-preferred subset of Y that has size less than or equal to a .

Y	$q(Y)$	$C^h(Y)$
$\{x, \hat{x}, y, z, \hat{z}\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{x, \hat{x}, y, z\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{x, \hat{x}, z, \hat{z}\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{\hat{x}, y, z, \hat{z}\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{x, \hat{x}, z\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{\hat{x}, z, \hat{z}\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{\hat{x}, y, z\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{\hat{x}, z\}$	(1, 0, 0, 1)	$\{\hat{x}, z\}$
$\{x, \hat{x}, y, \hat{z}\}$	(0, 1, 1, 0)	$\{x, \hat{z}\}$
$\{x, y, z, \hat{z}\}$	(0, 1, 1, 0)	$\{x, \hat{z}\}$
$\{x, \hat{x}, \hat{z}\}$	(0, 1, 1, 0)	$\{x, \hat{z}\}$
$\{x, y, \hat{z}\}$	(0, 1, 1, 0)	$\{x, \hat{z}\}$
$\{x, z, \hat{z}\}$	(0, 1, 1, 0)	$\{x, \hat{z}\}$
$\{x, \hat{z}\}$	(0, 1, 1, 0)	$\{x, \hat{z}\}$
$\{\hat{x}, y, \hat{z}\}$	(0, 1, 1, 0)	$\{y, \hat{z}\}$
$\{y, z, \hat{z}\}$	(0, 1, 1, 0)	$\{y, \hat{z}\}$
$\{y, \hat{z}\}$	(0, 1, 1, 0)	$\{y, \hat{z}\}$
$\{x, \hat{x}, y\}$	(1, 1, 0, 0)	$\{\hat{x}, y\}$
$\{\hat{x}, y\}$	(1, 1, 0, 0)	$\{\hat{x}, y\}$
$\{x, y, z\}$	(0, 2, 0, 0)	$\{x, y\}$
$\{x, y\}$	(0, 2, 0, 0)	$\{x, y\}$
$\{y, z\}$	(0, 1, 0, 1)	$\{y, z\}$
$\{\hat{x}, \hat{z}\}$	(1, 0, 1, 0)	$\{\hat{x}, \hat{z}\}$
$\{x, z\}$	(0, 1, 0, 1)	$\{x, z\}$
$\{y\}$	(0, 1, 0, 0)	$\{y\}$
$\{z, \hat{z}\}$	(0, 0, 1, 0)	$\{\hat{z}\}$
$\{\hat{z}\}$	(0, 0, 1, 0)	$\{\hat{z}\}$
$\{x, \hat{x}\}$	(1, 0, 0, 0)	$\{\hat{x}\}$
$\{\hat{x}\}$	(1, 0, 0, 0)	$\{\hat{x}\}$
$\{x\}$	(0, 1, 0, 0)	$\{x\}$
$\{z\}$	(0, 0, 0, 1)	$\{z\}$
\emptyset	(0, 0, 0, 0)	\emptyset

TABLE C1—THE VALUE OF THE ALLOTMENT FUNCTION AND CHOICE FUNCTION OF h FOR EVERY POSSIBLE SET OF CONTRACTS AVAILABLE TO h .

capacity constrained when $c(\mathbf{x})$ is available, and so we have nothing to show. If $|c(\mathbf{x})| = 3$, there are four cases to consider:

$c(\mathbf{x}) = \{x, y, z\}$. In this case, division 4—and only division 4—is capacity constrained under $c(\mathbf{x})$ at 0. Moreover, since \mathbf{y} is observable and $c(\mathbf{x}) \subsetneq c(\mathbf{y})$, we must have that $c(\mathbf{y}) = \{x, y, z, \hat{z}\}$, under which division 4 is still capacity constrained at 0.

$c(\mathbf{x}) = \{\hat{x}, y, \hat{z}\}$. In this case, division 1—and only division 1—is capacity constrained under $c(\mathbf{x})$ at 0. Moreover, since \mathbf{y} is observable and $c(\mathbf{x}) \subsetneq c(\mathbf{y})$, we must have that $c(\mathbf{y}) = \{\hat{x}, y, \hat{z}, x\}$, under which division 1 is still capacity constrained at 0.

$c(\mathbf{x}) = \{x, y, \hat{z}\}$. In this case, $q(c(\mathbf{x})) = (0, 1, 1, 0)$. Moreover, since \mathbf{y} is observable and $c(\mathbf{x}) \subsetneq c(\mathbf{y})$, we must have that $c(\mathbf{y}) = \{x, y, z, \hat{z}\}$; but then we also have $q(c(\mathbf{y})) = (0, 1, 1, 0)$.

$c(\mathbf{x}) = \{\hat{x}, y, z\}$. In this case, there does not exist an observable offer process \mathbf{y} such that $c(\mathbf{x}) \subsetneq c(\mathbf{y})$.

D. MATCHING WITH DISTRIBUTIONAL CONSTRAINTS

In this appendix, we explain how our framework nests the matching with regional caps model of Kamada and Kojima (2015, 2017).

D1. The Kamada and Kojima (2015, 2017) Model of Matching with Distributional Constraints

First, we must note one small, potentially confusing point of terminology: In the Kamada and Kojima (2015, 2017) framework, hospitals are partitioned into regions, and there are distributional constraints (regional caps) that restrict the number of doctors that can be assigned to each region. In our framework, by contrast, hospitals are the top-level institutions, and constraints within hospitals determine distributions across divisions. Thus, the divisions in our framework correspond to Kamada and Kojima’s (2015, 2017) *hospitals*, while our hospitals correspond to Kamada and Kojima’s (2015, 2017) *regions*.

In the setting of Kamada and Kojima (2015, 2017), doctors have strict preferences over the jobs they could have—represented in our context by contracts

$$(d, h, s) \in X = \bigcup_{h \in H} D \times \{h\} \times S^h$$

or, equivalently, by hospital–division pairs

$$(h, s) \in J \equiv \bigcup_{h \in H} \{(h, s) : s \in S^h\}.$$
^{6,7}

The preferences of doctor d over contracts in X_d naturally correspond to preferences over hospital–division pairs: If $(d, h, s) \succ_d (d, g, t)$ for some hospitals $h, g \in H$, some division $s \in S^h$, and some division $t \in S^g$, then we write $(h, s) \succ_d (g, t)$. Similarly, if $(d, h, s) \succ_d \emptyset$ for some hospital $h \in H$ and some division $s \in S^h$, then we write $(h, s) \succ_d \emptyset$. Finally, if $\emptyset \succ_d (d, h, s)$ for some hospital $h \in H$ and some division $s \in S^h$, then we write $\emptyset \succ_d (h, s)$.

Each hospital–division pair $j \in J$ has *responsive preferences* with respect to some strict ranking \succ_j of $D \cup \{\emptyset\}$ and a fixed *capacity* \bar{q}^j . That is, from any given set of contracts Y , the choice

⁶In the Kamada and Kojima (2015, 2017) model, doctors’ preferences could be expressed as rankings over just divisions, as Kamada and Kojima (2015, 2017) took divisions as primitives. In our context, doctors’ preferences must be expressed over hospital–division pairs (h, s) , as divisions in our setup only exist when associated with a hospitals.

⁷Here, unlike in the main text, we explicitly note the dependence of the set of divisions S on the underlying hospital, as we need to consider doctors’ preferences over hospital–division pairs.

function C^j of the hospital–division pair $j = (h, s)$ chooses the contracts in

$$Y_j \equiv \left\{ (d, \hat{h}, \hat{s}) \in Y : j = (\hat{h}, \hat{s}) = (h, s) \right\}$$

associated to the \bar{q}^j most highly-ranked doctors according to \succ_j ; if there are fewer than \bar{q}^j contracts in Y_j associated with doctors that j ranks more highly than the outside option \emptyset , then choice function C^j of j chooses all such contracts. Additionally, each hospital h has an *overall capacity* \bar{Q}^h .

A *matching* is a mapping μ that assigns doctors to hospital–division pairs, i.e., a mapping μ such that

- 1) $\mu(d) \in J \cup \{\emptyset\}$ for all $d \in D$,
- 2) $\mu(j) \subseteq D$ for all $j \in H$, and
- 3) for all $d \in D$ and $h \in H$, we have that $d \in \mu(j)$ if and only if $\mu(d) = j$.

A matching μ is

- *individually rational for doctors* if, for all $d \in D$, we have $\mu(d) \succ_d \emptyset$;
- *individually rational for hospital divisions* if, for all hospital–division pairs $j \in J$, we have $d \succ_j \emptyset$ whenever $d \in \mu(j)$;
- *feasible* if $|\mu(j)| \leq \bar{q}^j$ for each $j \in J$ and $\sum_{s \in S^h} |\mu((h, s))| \leq \bar{Q}^h$ for each $h \in H$; and
- *blocked by* $(d, j) \in D \times J$ if $j \succ_d \mu(d)$, $d \succ_j \emptyset$, and either $|\mu(j)| < \bar{q}^j$ or $d \succ_j e$ for some $e \in \mu(j)$.

At the aggregate level, the hospital h has preferences over capacity distributions across divisions. Formally, there is a weak ordering \succeq_h over the set of *distribution vectors*

$$\mathcal{W}^h \equiv \{(w_s)_{s \in S^h} : w_s \in \mathbb{Z}_{\geq 0}\}.$$

Given \succeq_h , a *capacity allocation rule* is a mapping $p : \mathcal{W}^h \rightarrow \mathcal{W}^h$ such that for all $w \in \mathcal{W}^h$,

$$p(w) \in \max_{\succeq_h} \{w' : w' \leq w\}.$$
^{8,9}

Kamada and Kojima (2015, 2017) imposed the following assumptions on the capacity allocation rule p :

- 1) If $w, w' \in \mathcal{W}^h$ are such that $p(w) \leq w' \leq w$, then $p(w') = p(w)$.
- 2) For all $w \in \mathcal{W}^h$ and all $s \in S^h$, $[p(w)]_s \leq \bar{q}^{(h,s)}$.
- 3) For all $w \in \mathcal{W}^h$, $\sum_{s \in S^h} [p(w)]_s \leq \bar{Q}^h$.
- 4) For all $w \in \mathcal{W}^h$, if there is an $s \in S^h$ such that $[p(w)]_s < \min\{w_s, \bar{q}^{(h,s)}\}$, then $\sum_{t \in S^h} [p(w)]_t = \bar{Q}^h$.
- 5) For all $w, w' \in \mathcal{W}^h$ and $s \in S^h$ such that $w \leq w'$ and $[p(w)]_s < [p(w')]_s$, we have $[p(w)]_s = w_s$.

In the sequel, we assume the preceding conditions, and refer to them as *Conditions 1–5 of Kamada and Kojima (2015, 2017)*.

Kamada and Kojima (2015, 2017) also introduced the following stability concept, *stability under distributional constraints*.

⁸Here, we suppress the dependence of p on the hospital h for notational simplicity.

⁹Here, by \max_{\succeq_h} , we mean the maximum with respect to the ordering \succeq_h .

DEFINITION 1: A matching μ is stable under distributional constraints, if it is feasible and individually rational for doctors and hospitals, and whenever $(d, (h, s))$ is a blocking pair the following three conditions hold:

- 1) The hospital h is capacity constrained, i.e., $\sum_{\hat{s} \in S^h} |\mu((h, \hat{s}))| = \bar{Q}^h$.
- 2) The hospital–division pair (h, s) prefers all of its doctors under μ to d , i.e., $d' \succ_{(h,s)} d$ for all $d' \in \mu((h, s))$.
- 3) Either
 - a) doctor d is not employed at hospital h under μ , i.e., $d \notin \cup_{s \in S^h} \mu((h, s))$, or
 - b) hospital h prefers its distribution vector under μ to the one that would arise if d were to switch to j , that is, $(|\mu((h, \hat{s}))|)_{\hat{s} \in S^h} \succeq_h v$, where

$$v_{\hat{s}} = \begin{cases} |\mu((h, \hat{s}))| + 1 & \hat{s} = s \\ |\mu((h, \hat{s}))| - 1 & (h, \hat{s}) = \mu(d) \\ |\mu((h, \hat{s}))| & \text{otherwise.} \end{cases}$$

Definition 1 rules out blocks $(d, (h, s))$ in which the hospital h is capacity constrained, the division s only benefits if it adds d as a new doctor, and either d is employed at a different hospital pre-block or the hospital h prefers its distribution vector pre-block to its distribution vector post-block.

D2. Embedding the Kamada–Kojima (2015, 2017) Model within Our Framework

We now show how to embed the model of Kamada and Kojima (2015, 2017) into our model of matching with flexible allotments. For notational simplicity, we focus on a single hospital h , and return to suppressing the notation for h wherever doing so will not introduce confusion. Now, for each doctor d and division $s \in S^h$, there is just one contract, denoted $(d, s) = (d, h, s)$, under which d is employed at division s (of h). Thus, the set of contracts can be reduced to $X = \cup_{s \in S^h} \{(d, s) : d \in D\}$.

We assume that each division $s \in S^h$ has a strict ranking \succ_s of contracts in the set

$$\{(d, s) \in X : d \in D\} \cup \{\emptyset\}$$

such that

- 1) $(d, s) \succ_s (d', s)$ if and only if $d \succ_{(h,s)} d'$,
- 2) $(d, s) \succ_s \emptyset$ if and only if $d \succ_{(h,s)} \emptyset$, and
- 3) $\emptyset \succ_s (d, s')$ for all $s' \neq s$ and $d \in D$.¹⁰

For any allotment a , the extended choice function $C^s(\cdot; a)$ of each division s is assumed to be responsive with respect to the order \succ_s with capacity a .¹¹ Next, we use p to define an allotment function q by setting, for each $Y \subseteq X$,

$$(D1) \quad q(Y) = p(\varpi(Y)),$$

¹⁰Note that different divisions never have acceptable contracts in common.

¹¹We could instead use the capacity $\min\{a, \bar{q}^{(h,s)}\}$, which bounds the number of accepted doctors at $\bar{q}^{(h,s)}$. This is not formally necessary in our construction, however, as we impose the division cap $\bar{q}^{(h,s)}$ via the allotment function.

where we take

$$(D2) \quad \varpi(Y) \equiv (|\{y \in Y_h : y \succ_{(h,s)} \emptyset\}|)_{s \in S}$$

to be the vector that counts the number of contracts in Y_h that are acceptable to each division s .

With our specifications of the extended choice functions C^s and the allotment function q , the choice function C^h implements the choices made by the hospital–division pairs (h, s) under the choice functions $C^{(h,s)}$ and the constraints imposed by p under the cumulative offer mechanism.

We now prove that the extended choice functions C^s , combined with the allotment function q , do indeed make the induced hospital choice function C^h a multi-division choice function with flexible allotments.

CLAIM 7: *The hospital choice function C^h induced by $(C^s)_{s \in S}$ and q (under the choice procedure defined in Section III) is a multi-division choice function with flexible allotments. That is:*

- For any fixed allotment, each division’s extended choice function $C^s(\cdot; a)$ is substitutable and size monotonic, and satisfies the irrelevance of rejected contracts condition. Moreover, each division’s extended choice function C^s is monotonic with respect to the allotment and conditionally acceptant.
- The allotment function q does not depend on irrelevant contracts, does not observably grant excess positions, is single-peaked across observable offer processes, and is monotone in aggregate across observable offer processes.

PROOF:

As $C^s(\cdot; a)$ is responsive, it is immediate that it satisfies the classical substitutability, size monotonicity, and irrelevance of rejected contracts conditions. Likewise, it is immediate that each division’s extended choice function C^s is monotonic with respect to the allotment and conditionally acceptant.

Now, we show the claimed properties of the allotment function q ; as we do so, we sometimes abuse notation slightly by writing

$$p_s(w) \equiv (p(w))_s \quad \text{and} \quad \varpi_s(Y) \equiv (\varpi(Y))_s.$$

1) **The allotment function q does not depend on irrelevant contracts:** We consider a set of contracts $Y \subseteq X$ and suppose that $z \in R^h(Y)$. We first argue that

$$(D3) \quad q(Y) = p(\varpi(Y)) \leq \varpi(Y \setminus \{z\}).$$

First, we recall that the contract z is associated to a unique division—that is, $z = (d, s)$ for some division s . Thus, if $z = (d, s) \in R^h(Y)$, we must have either:

- $\emptyset \succ_{(h,s)} d$ (d is unacceptable to s),
- $d \succ_{(h,s)} \emptyset$ and $p_s(\varpi(Y)) = \bar{q}^{(h,s)}$ (d is acceptable to s , but s is at maximum-possible capacity), or
- $d \succ_{(h,s)} \emptyset$ and $p_s(\varpi(Y)) < \bar{q}^{(h,s)}$ (d is acceptable to s , but the allotment rule constrains s below its maximum-possible capacity).

In the first case, we must have $\varpi_t(Y) = \varpi_t(Y \setminus \{z\})$ for all $t \in S$ (recall (D2)); hence, (D3) is immediate, as $p_t(\varpi_t(Y)) \leq \varpi_t(Y)$ by construction.

In the second and third cases, we have that $q^s(Y) \leq \varpi_s(Y) - 1$ given that $z \in R^h(Y)$ and $d \succ_{(h,s)} \emptyset$; as $p_s(\varpi(Y)) = q^s(Y)$, this implies that

$$(D4) \quad p_s(\varpi(Y)) \leq \varpi_s(Y) - 1.$$

Meanwhile, we have

$$(D5) \quad \varpi_s(Y \setminus \{z\}) = \varpi_s(Y) - 1,$$

as $|\{y \in Y_h \setminus \{z\} : y \succ_{(h,s)} \emptyset\}| = |\{y \in Y_h : y \succ_{(h,s)} \emptyset\}| - 1$. Combining (D4) with (D5) shows that

$$(D6) \quad p_s(\varpi(Y)) \leq \varpi_s(Y \setminus \{z\}).$$

Meanwhile, as z is associated to s , it is unacceptable to all divisions $t \neq s$; this implies that $\varpi_t(Y \setminus \{z\}) = \varpi_t(Y)$ for all such t . As $p_t(\varpi(Y)) \leq \varpi_t(Y)$ for all $t \in S$, we then have

$$(D7) \quad p_t(\varpi(Y)) \leq \varpi_t(Y \setminus \{z\}) \quad \text{for all divisions } t \neq s.$$

Combining (D6) with (D7), we find that

$$p_t(\varpi(Y)) \leq \varpi_t(Y \setminus \{z\})$$

for *all* divisions $t \in S$ —exactly (D3).

Now, having proven (D3), we note that $\varpi_t(Y \setminus \{z\}) \leq \varpi_t(Y)$ mechanically, so that Condition 1 of Kamada and Kojima (2015, 2017) implies that $p(\varpi(Y)) = p(\varpi(Y \setminus \{z\}))$. Thus, we have $q(Y \setminus \{z\}) = p(\varpi(Y \setminus \{z\})) = p(\varpi(Y)) = q(Y)$; this implies the claim.

- 2) **The allotment function q does not observably grant excess positions:** Let $\mathbf{x} = (x^1, \dots, x^M)$ be an observable offer process, and assume that there is a division s such that

$$(D8) \quad q^s(\mathbf{c}(\mathbf{x})) > \min\{|\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}) : d \succ_{(h,s)} \emptyset\}|, \bar{q}^{(h,s)}\}.$$

We assume without loss of generality that¹²

$$(D9) \quad \begin{aligned} &\text{for all pairs } (m, t) \text{ such that either } m < M \text{ or } m = M \text{ and } t < s, \\ &\text{we have } q^t(\mathbf{c}(\mathbf{x}^m)) \leq \min\{|\{(d, t) \in \mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^m) : d \succ_{(h,t)} \emptyset\}|, \bar{q}^{(h,t)}\}. \end{aligned}$$

Now, as $|\mathbf{c}(\mathbf{x}) \setminus \mathbf{c}(\mathbf{x}^{M-1})| = 1$, we must have $p_t(\varpi(\mathbf{c}(\mathbf{x}))) \leq p_t(\varpi(\mathbf{c}(\mathbf{x}^{M-1}))) + 1$ for all t . Moreover, by Condition 5 of Kamada and Kojima (2015, 2017), we have $p_t(\varpi(\mathbf{c}(\mathbf{x}))) = p_t(\varpi(\mathbf{c}(\mathbf{x}^{M-1}))) + 1$ only if $p_t(\varpi(\mathbf{c}(\mathbf{x}^{M-1}))) = \varpi_t(\mathbf{c}(\mathbf{x}^{M-1}))$.

Combining the preceding observations with (D9) (and recalling that $q(\cdot) = p(\varpi(\cdot))$ by (D1)), we see that for all pairs (m, t) such that either $m < M$ or $m = M$ and $t < s$, we must have either

$$(D10) \quad q^t(\mathbf{c}(\mathbf{x})) \leq q^t(\mathbf{c}(\mathbf{x}^{M-1}))$$

or

$$(D11) \quad \begin{aligned} q^t(\mathbf{c}(\mathbf{x})) &= q^t(\mathbf{c}(\mathbf{x}^{M-1})) + 1 \\ q^t(\mathbf{c}(\mathbf{x}^{M-1})) &= |\{(d, t) \in \mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{M-1}) : d \succ_{(h,t)} \emptyset\}| < \bar{q}^{(h,t)} \\ &\quad \mathbf{d}(x^M) \succ_{(h,t)} \emptyset \\ &\quad x^M \in \mathbf{c}_{\rightsquigarrow t}(\mathbf{x}) \setminus \mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{M-1}). \end{aligned}$$

¹²If (D9) did not hold for the claimed pairs (m, t) , then we could shorten the offer process \mathbf{x} to \mathbf{x}^m or replace s with t .

Now, we prove via induction that

$$(D12) \quad C^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{M-1}); q^t(\mathbf{c}(\mathbf{x}^{M-1}))) \subseteq \mathbf{c}_{\rightsquigarrow t}(\mathbf{x}) \text{ for all } t \leq s.$$

First, we note that (D12) is immediately satisfied for the base case of $t = 1$, as

$$\mathbf{c}_{\rightsquigarrow 1}(\mathbf{x}^{M-1}) = \mathbf{c}(\mathbf{x}^{M-1}) \subseteq \mathbf{c}(\mathbf{x}) = \mathbf{c}_{\rightsquigarrow 1}(\mathbf{x}).$$

We suppose that (D12) holds for all $t' < t \leq s$, and fix some $y \in C^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{M-1}); q^t(\mathbf{c}(\mathbf{x}^{M-1})))$. Since $y \in C^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{M-1}); q^t(\mathbf{c}(\mathbf{x}^{M-1})))$, under the choice procedure defining C^h , for any $t' < t$, we must have that the contract y is not among the $q^{t'}(\mathbf{c}(\mathbf{x}^{M-1}))$ most preferred contracts in $\{(d, t') \in \mathbf{c}_{\rightsquigarrow t'}(\mathbf{x}^{M-1}) : d \succ_{(h, t')} \emptyset\}$.

If $q^{t'}(\mathbf{c}(\mathbf{x})) \leq q^{t'}(\mathbf{c}(\mathbf{x}^{M-1}))$, then (D12) in the t' case immediately implies that y is not among the $q^{t'}(\mathbf{c}(\mathbf{x}))$ most preferred contracts in $\{(d, t') \in \mathbf{c}_{\rightsquigarrow t'}(\mathbf{x}) : d \succ_{(h, t')} \emptyset\}$.

Otherwise, if $q^{t'}(\mathbf{c}(\mathbf{x})) = q^{t'}(\mathbf{c}(\mathbf{x}^{M-1})) + 1$, we must have that

$$q^{t'}(\mathbf{c}(\mathbf{x}^{M-1})) = |\{(d, t') \in \mathbf{c}_{\rightsquigarrow t'}(\mathbf{x}^{M-1}) : d \succ_{(h, t')} \emptyset\}| < \bar{q}^{(h, t')},$$

recalling (D11). As y is not chosen by t' when $\mathbf{c}(\mathbf{x}^{M-1})$ is available to h , the preceding observation implies that either $\emptyset \succ_{(h, t')} d(y)$ or $(d(y), t') \notin \mathbf{c}(\mathbf{x}^{M-1})$. In the former case, it is immediate that $d(y) \notin d(C^{t'}(\mathbf{c}_{\rightsquigarrow t'}(\mathbf{x}); q^{t'}(\mathbf{c}(\mathbf{x}))))$. In the latter case, $y \in C^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{M-1}); q^t(\mathbf{c}(\mathbf{x}^{M-1})))$ implies $d(y) \neq d(x^M)$ and thus $(d(y), t') \notin \mathbf{c}(\mathbf{x})$.

Hence, no matter whether $q^{t'}(\mathbf{c}(\mathbf{x})) \leq q^{t'}(\mathbf{c}(\mathbf{x}^{M-1}))$ or $q^{t'}(\mathbf{c}(\mathbf{x})) = q^{t'}(\mathbf{c}(\mathbf{x}^{M-1})) + 1$, we obtain that $d(y) \notin d(C^{t'}(\mathbf{c}_{\rightsquigarrow t'}(\mathbf{x}); q^{t'}(\mathbf{c}(\mathbf{x}))))$. Since y was arbitrary, this shows that $C^t(\mathbf{c}_{\rightsquigarrow t}(\mathbf{x}^{M-1}); q^t(\mathbf{c}(\mathbf{x}^{M-1}))) \subseteq \mathbf{c}_{\rightsquigarrow t}(\mathbf{x})$ if $C^{t'}(\mathbf{c}_{\rightsquigarrow t'}(\mathbf{x}^{M-1}); q^{t'}(\mathbf{c}(\mathbf{x}^{M-1}))) \subseteq \mathbf{c}_{\rightsquigarrow t'}(\mathbf{x})$ for all $t' < t$; this completes the proof of (D12).

To complete the proof that q does not observably grant excess positions, we now derive a contradiction to (D8). Note again from (D9) that

$$q^s(\mathbf{c}(\mathbf{x}^{M-1})) \leq \min\{|\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}) : d \succ_{(h, s)} \emptyset\}|, \bar{q}^{(h, s)}\};$$

this inequality implies that $|C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1})))| = q^s(\mathbf{c}(\mathbf{x}^{M-1}))$. There are two cases to consider:

- We consider first the case in which $q^s(\mathbf{c}(\mathbf{x})) \leq q^s(\mathbf{c}(\mathbf{x}^{M-1}))$. Given (D12), we have $C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1}))) \subseteq \mathbf{c}_{\rightsquigarrow s}(\mathbf{x})$, and thus

$$|C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1})))| \leq |\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}) : d \succ_{(h, s)} \emptyset\}|.$$

Since $|C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1})))| = q^s(\mathbf{c}(\mathbf{x}^{M-1}))$ and $q^s(\mathbf{c}(\mathbf{x})) \leq q^s(\mathbf{c}(\mathbf{x}^{M-1}))$, we obtain a contradiction to (D8).

- We consider second the case in which $q^s(\mathbf{c}(\mathbf{x})) > q^s(\mathbf{c}(\mathbf{x}^{M-1}))$. Note that, here again, (D11) must hold. Since we then have $q^s(\mathbf{c}(\mathbf{x}^{M-1})) = |\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}) : d \succ_{(h, s)} \emptyset\}|$ and $|\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}) : d \succ_{(h, s)} \emptyset\}| < \bar{q}^{(h, s)}$, we see that $C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1}))) = \{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}) : d \succ_{(h, s)} \emptyset\}$. Since $C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1}))) \subseteq \mathbf{c}_{\rightsquigarrow s}(\mathbf{x})$ and $x^M \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}) \setminus C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}); q^s(\mathbf{c}(\mathbf{x}^{M-1})))$, we have that

$$|\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}) : d \succ_{(h, s)} \emptyset\}| = |\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}) : d \succ_{(h, s)} \emptyset\}| + 1.$$

Since, again by (D11), we have that $|\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}^{M-1}) : d \succ_{(h, s)} \emptyset\}| = q^s(\mathbf{c}(\mathbf{x}^{M-1}))$

and $q^s(\mathbf{c}(\mathbf{x})) = q^s(\mathbf{c}(\mathbf{x}^{M-1})) + 1$, we obtain that $|\{(d, s) \in \mathbf{c}_{\rightsquigarrow s}(\mathbf{x}) : d \succ_{(h,s)} \emptyset\}| = q^s(\mathbf{c}(\mathbf{x}))$, again contradicting our earlier assumption (D8).

- 3) **The allotment function q is single-peaked across observable offer processes:** We consider observable offer processes \mathbf{x} and \mathbf{y} such that $\mathbf{c}(\mathbf{x}) \subseteq \mathbf{c}(\mathbf{y})$. Let s be a division such that $|C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); \infty)| > q^s(\mathbf{c}(\mathbf{x}))$. As $\varpi_s(\mathbf{c}(\mathbf{x})) \geq |C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); \infty)|$ automatically, we see that $\varpi_s(\mathbf{c}(\mathbf{x})) > q^s(\mathbf{c}(\mathbf{x}))$. As $q(\mathbf{c}(\mathbf{x})) = p(\varpi(\mathbf{c}(\mathbf{x})))$ by construction (recall (D1)) and $\varpi(\mathbf{c}(\mathbf{y})) \geq \varpi(\mathbf{c}(\mathbf{x}))$ automatically, Condition 5 of Kamada and Kojima (2015, 2017) then implies that $p_s(\varpi(\mathbf{c}(\mathbf{y}))) \leq p_s(\varpi(\mathbf{c}(\mathbf{x})))$, so that $q^s(\mathbf{c}(\mathbf{y})) \leq q^s(\mathbf{c}(\mathbf{x}))$, as desired.
- 4) **The allotment function q is monotone in aggregate across observable offer processes:** We consider observable offer processes \mathbf{x} and \mathbf{y} such that $\mathbf{c}(\mathbf{x}) \subseteq \mathbf{c}(\mathbf{y})$. If we have

$$(D13) \quad \sum_{s \in S} q^s(\mathbf{c}(\mathbf{x})) > \sum_{s \in S} q^s(\mathbf{c}(\mathbf{y})),$$

then it must be the case that $\sum_{s \in S} q^s(\mathbf{c}(\mathbf{y})) < \bar{Q}^h$, and so, by Condition 4 of Kamada and Kojima (2015, 2017), we have that $q^s(\mathbf{c}(\mathbf{y})) = p_s(\varpi(\mathbf{c}(\mathbf{y}))) = \min\{\varpi_s(\mathbf{c}(\mathbf{y})), \bar{q}^{(h,s)}\}$ for all s (as, by the definition of the capacity allocation rule, $q^s(\mathbf{c}(\mathbf{y})) \leq \varpi_s(\mathbf{c}(\mathbf{y}))$), and, by Condition 2 of Kamada and Kojima (2015, 2017), $q^s(\mathbf{c}(\mathbf{y})) \leq \bar{q}^{(h,s)}$). But, since $\mathbf{c}(\mathbf{x}) \subseteq \mathbf{c}(\mathbf{y})$, we have that $\min\{\varpi_s(\mathbf{c}(\mathbf{x})), \bar{q}^{(h,s)}\} \leq \min\{\varpi_s(\mathbf{c}(\mathbf{y})), \bar{q}^{(h,s)}\}$ for all s . Thus, by Condition 4 of Kamada and Kojima (2015, 2017),

$$q^s(\mathbf{c}(\mathbf{x})) = p_s(\varpi(\mathbf{c}(\mathbf{x}))) \leq \min\{\varpi_s(\mathbf{c}(\mathbf{x})), \bar{q}^{(h,s)}\} \leq \min\{\varpi_s(\mathbf{c}(\mathbf{y})), \bar{q}^{(h,s)}\} = q^s(\mathbf{c}(\mathbf{y})).$$

But then, summing over all s , we have that

$$\sum_{s \in S} q^s(\mathbf{c}(\mathbf{x})) \leq \sum_{s \in S} q^s(\mathbf{c}(\mathbf{y})),$$

contradicting (D13).

D3. Strategy-Proofness of Cumulative Offer Mechanisms in the Kamada–Kojima (2015, 2017) Setting

With the embedding described in Appendix D2, strategy-proofness of cumulative offer mechanisms in the Kamada and Kojima (2015, 2017) context follows directly from our main results.

PROPOSITION 1: *Suppose that each hospital has a choice function constructed as in the model of Kamada and Kojima (2015, 2017) (as described in Appendix D1). Then any cumulative offer mechanism is strategy-proof.*

PROOF:

This follows immediately upon combining Claim 7 with Corollary 1.

D4. Stability of Cumulative Offer Mechanism Outcomes Under Distributional Constraints

Finally, we show that cumulative offer mechanism outcomes in our context correspond to matchings that are stable under distributional constraints.

For each feasible set of contracts $Y \subseteq X$, we define a matching μ^Y by setting

$$\mu^Y(d) = \begin{cases} (h, s) & (d, h, s) \in Y \\ \emptyset & \text{otherwise,} \end{cases}$$

$$\mu^Y((h, s)) = \{d : (d, h, s) \in Y\}.$$

PROPOSITION 2: *Suppose that each hospital has a choice function constructed as in the model of Kamada and Kojima (2015, 2017) (as described in Appendix D1). Then if A is the outcome of a cumulative offer mechanism, the matching μ^A is stable under distributional constraints.*

PROOF:

We fix the preferences of the doctors and hospital–division pairs as well as the capacity allocation rules of hospitals, and use the construction described in Appendix D2 to formulate the associated choice functions and preferences over contracts, as well as the allotment functions. We let $\mathbf{x} = (x^1, \dots, x^M)$ be the sequence of contracts proposed under any cumulative offer mechanism given those preferences.¹³ With this setup, we have

$$A = \bigcup_{h \in H} C^h(\mathbf{c}(\mathbf{x})).$$

It is immediate that μ^A is

- feasible and
- individually rational for both doctors and hospital divisions.

Now, suppose that μ^A is blocked by $(d, (h, s))$, i.e., $(h, s) \succ_d \mu^A(d)$ and $d \succ_{(h,s)} \emptyset$.

We argue first that $\hat{d} \succ_{(h,s)} d$ for all $\hat{d} \in \mu^A(s)$ (i.e., Condition 2 of Definition 1). Since $(h, s) \succ_d \mu^A(d)$, we must have that $x = (d, h, s)$ was proposed at some step of the cumulative offer process corresponding to \mathbf{x} . Since $(d, h, s) \notin A$, there has to exist an \hat{M} such that, letting $\hat{\mathbf{x}} \equiv (x^1, \dots, x^{\hat{M}})$, we have that $(d, h, s) \in R^s(\mathbf{c}_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(\mathbf{c}(\hat{\mathbf{x}})))$. By (B1) of Claim 1, we must have that $(d, h, s) \in R^s(\mathbf{f}_{\rightsquigarrow s}(\hat{\mathbf{x}}); q^s(\mathbf{c}(\hat{\mathbf{x}})))$. By (B2) of Claim 1, we must then have that

$$(D14) \quad (d, h, s) \in R^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x}))).$$

Now, if $\hat{d} \in \mu^A((h, s))$, then $(\hat{d}, h, s) \in C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x})))$. Moreover, by (B1) of Claim 1, we must have that $C^s(\mathbf{c}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x}))) = C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x})))$, and so

$$(D15) \quad (\hat{d}, h, s) \in C^s(\mathbf{f}_{\rightsquigarrow s}(\mathbf{x}); q^s(\mathbf{c}(\mathbf{x}))).$$

Together, (D14) and (D15) imply that $\hat{d} \succ_{(h,s)} d$ for all $\hat{d} \in \mu^A(s)$, as desired.

We argue second that h is capacity constrained (Condition 1 of Definition 1). We have just shown that $\hat{d} \succ_{(h,s)} d$ for all $\hat{d} \in \mu^A(s)$. Nevertheless, $(d, (h, s))$ blocks μ ; hence, it must be the case that $|\mu^A((h, s))| < \bar{q}^{(h,s)}$.¹⁴ Moreover, as $\mu(d) \neq (h, s)$ under μ even though (d, h, s) is both

- in $\mathbf{c}(\mathbf{x})$ and
- acceptable to (h, s) ,

we must have $|\mu^A((h, s))| < \varpi_s(\mathbf{c}(\mathbf{x}))$. Thus, we have

$$p_s(\varpi(\mathbf{c}(\mathbf{x}))) = q^s(\mathbf{c}(\mathbf{x})) = |\mu^A((h, s))| < \min\{\varpi_s(\mathbf{c}(\mathbf{x})), \bar{q}^{(h,s)}\};$$

hence, Condition 4 of Kamada and Kojima (2015, 2017) implies that hospital h is capacity constrained, as desired.

¹³By Proposition 1 of Hatfield et al. (2016), all cumulative offer mechanisms in this context are outcome-equivalent.

¹⁴If we had $|\mu^A((h, s))| = \bar{q}^{(h,s)}$ then $(d, (h, s))$ could not block μ , as under μ the hospital–division pair (h, s) would be assigned $\bar{q}^{(h,s)}$ doctors it prefers to d .

Finally, we show Condition 3 of Definition 1. Suppose that d is employed at h under μ (Case 3b of Condition 3), i.e., there exists a division $\hat{s} \in S^h$ such that $\mu(d) = (h, \hat{s})$. By construction, we have

$$p(\varpi(\mathbf{c}(\mathbf{x}))) = q(\mathbf{c}(\mathbf{x})) = (|\mu^A((h, t))|)_{t \in S^h}.$$

Let v be obtained by setting $v_s = p_s(\varpi(\mathbf{c}(\mathbf{x}))) + 1$, $v_{\hat{s}} = p_{\hat{s}}(\varpi(\mathbf{c}(\mathbf{x}))) - 1$, and $v_t = p_t(\varpi(\mathbf{c}(\mathbf{x})))$ for all $t \in S^h \setminus \{s, \hat{s}\}$. As $(d, h, s) \in \mathbf{c}(\mathbf{x})$ and $(d, h, s) \notin \mu^A((h, s))$, we must have $v_s < \varpi_s(\mathbf{c}(\mathbf{x}))$, so that $v \leq \varpi(\mathbf{c}(\mathbf{x}))$. But then, $v \in \{w' : w' \leq \varpi(\mathbf{c}(\mathbf{x}))\}$. Thus, we must have

$$(|\mu^A((h, t))|)_{t \in S^h} = p(\varpi(\mathbf{c}(\mathbf{x}))) \succeq_h v,$$

as when allocating capacity under p , the hospital h could have chosen the distribution vector v but instead chose the distribution vector corresponding to μ^A . Thus, we find that $(|\mu^A((h, t))|)_{t \in S^h} \succeq_h v$, as required by Definition 1.

*

REFERENCES

- Hatfield, John William and Scott Duke Kominers**, “Hidden Substitutes,” 2016. Working paper.
- , – , and **Alexander Westkamp**, “Stability, Strategy-Proofness, and Cumulative Offer Mechanisms,” 2016. Working paper.
- Kamada, Yuichiro and Fuhito Kojima**, “Efficient Matching under Distributional Constraints: Theory and Applications,” *American Economic Review*, 2015, 105 (1), 67–99.
- and – , “Recent Developments in Matching with Constraints,” *American Economic Review Papers & Proceedings*, 2017.